



IFSSH Scientific Committee on Carpal Instability

Part 1: Definition and Investigations

Chair: Max Haerle (Germany)

**Committee: Abhijeet Wahegaonkar (India)
Marc Garcia-Elias (Spain)
Gregory Bain (Australia)
Riccardo Luchetti (Italy)**

Report submitted May 2016

Part 1: Carpal Instability - Definition and Investigations

Introduction

The wrist is a load bearing articulation able to resist both compressive and torsional loads without yielding. For this to happen, there is a need for: 1) a coordinated action of the muscles crossing the joint, 2) smooth and normally tilted joint articular surfaces, and 3) a system of interdependent ligaments. In the presence of muscle imbalance, ligament insufficiency and/or bone deformity several patterns of wrist instability may appear. From this point of view, carpal instability may be described as the inability of the wrist to maintain a normal balance between the articulating surfaces under physiologic loads (dyskinetics) and/or movements (dyskinematics). When the wrist is not able to keep normal articular alignment when subjected to physiologic deforming forces, the wrist is said to be unstable.

Not long ago, carpal instabilities were only identified if the patient was referred to a knowledgeable wrist specialist. Nowadays, the diagnosis is often established in the emergency rooms of our hospitals. Certainly, the general understanding of these complex injuries has improved. Such an increased awareness, however, has not been followed by an increase in the quality of the results obtained. Indeed, the literature concerning the management of carpal instabilities still is rich in misconception justifying surgical techniques that have not yet proved the test of time. We are certainly improving, but we are not definitively there yet.

The natural history of carpal instabilities is still unclear. It is not unusual to find authors defending the notion that all ligament injuries will sooner or later evolve hopelessly into a symptomatic osteoarthritis. Now we know that ligament ruptures do not always generate instability, nor is all wrist osteoarthritis derived from an unstable joint. Furthermore, some malaligned, theoretically unstable wrists are well tolerated, if not completely asymptomatic. E.g. those produced by malunited fractures of the lower end of the radius or the dynamic midcarpal instability in patients with hyperlaxity of their joints. Others, such as scapho-lunate dissociation, may rapidly lead to degenerative arthritis and require early detection and treatment.

Classification

Carpal instability is difficult to classify. Many diverse clinical conditions may result in an unstable wrist for which different classifications have been suggested. Carpal instability has been classified based on seven parameters:

- 1) **Aetiology** (congenital, developmental, post-traumatic),
- 2) **Location of the initial injury** (extrinsic, intrinsic ligament, bone),
- 3) **Characteristics of the original injury** (partial vs. total; repairable vs non-repairable, reducible vs. non-reducible),

- 4) **Constancy of the radiological findings** (pre-dynamic, dynamic, static),
- 5) **Location of the predominant dysfunction** (dissociative scapholunate, dissociative luno-triquetral, non-dissociative radiocarpal, non-dissociative midcarpal),
- 6) **Chronicity of the dysfunction** (acute, sub-acute, chronic),
- 7) **Direction of the resultant malalignment** (DISI, VISI, ulnar translocation).

Unfortunately, none of these provide a comprehensive enough classification which is able to be used to decide a treatment. Algorithms of treatment based on a combination of the seven parameters have also been proposed. The ideal algorithm must include all possible forms of carpal instability, and be simple enough as to be easily remembered. Although imperfect, the analytical scheme proposed by Larsen and Associates in 1995 (Table 1) and the recently revised algorithm of treatment originally proposed by Garcia-Elias, Lluch and Stanley (2006) fulfil these criteria. Needless to say, there is a need for further refinement in this regards.

Table 1: A summary of classification types for carpal instability is presented below in a tabular form⁵

| Category I Chronicity | Category II Constancy | Category III Etiology | Category IV Location | Category V Direction | Category VI Pattern |
|---|--|---|---|--|---|
| Acute < 1 week (Maximum primary healing potential) | Predynamic Dynamic Static reducible | Congenital Traumatic Inflammatory | Radiocarpal Proximal Intercarpal Midcarpal | VISI rotation DISI rotation Ulnar translation | Carpal instability dissociative (CID) |
| Subacute 1-6 wks (some healing potential) | Static irreducible | Neoplastic Iatrogenic Miscellaneous | Distal intercarpal Carpometacarpal Specific bones | Carpal instability Dorsal translation Other | Carpal instability non- dissociative (CIND) |
| Chronic >6 wks (little healing potential) | | | | | Carpal instability complex (CIC) |
| | | | | | Carpal instability adaptive (CIA) |

Scapholunate Dissociation

Introduction

Injuries to the scapholunate ligament and the secondary restraints may lead to different degrees of scapholunate instability. Dependently this may lead to a considerable degree of wrist dysfunction, inability to work and interference with manual activities. If left untreated, it can lead to wrist osteoarthritis. Impairment of the scapholunate interosseous ligament (SLIL) in association with injury to the extrinsic ligaments is known to lead to rotatory subluxation of the scaphoid, dorsal intercalated segment instability (DISI) and finally scapholunate advanced collapse (SLAC).⁷

Anatomy

The SL ligament consists of three distinct structures: the two SL ligaments (palmar and dorsal) and the proximal fibrocartilaginous membrane.⁸

The **dorsal** SL ligament is located in the depth of the dorsal capsule and connects the dorsal aspects of the scaphoid and lunate bones. It is formed by thick and stout collection of fibers, slightly obliquely oriented, with a key role in SL stability. The dorsal component is a true ligament with transversely oriented collagen fibers, and is a primary restraint not only to distraction, but also to torsional and translational moments.

The **palmar** SL ligament, although considerably thinner, has important contributions to rotational stability of the SL joint.

The proximal **membranous** portion of the SLIL is histologically a fibrocartilaginous structure, and in isolation, contributes little to the restraint of the normal motion of the SL joint⁹. However, recent publications have also highlighted the major role of the secondary restraints, the dorsal intercarpal and radiocarpal ligaments in maintaining scapholunate stability.

In a study by Elsaidi et al¹⁰; the authors found that after sequential sectioning of volar ligaments and the scapholunate interosseous ligament, no scapholunate diastasis or excessive scaphoid flexion occurred. After dividing the dorsal intercarpal ligament, scapholunate instability occurred without carpal collapse. With detaching the dorsal radiocarpal ligament from the lunate, a dorsal intercalated scapholunate instability deformity ensued.

More recently, van Overstraeten et al¹¹ described an attachment between the dorsal wrist capsule, the dorsal part of the scapholunate interosseous ligament (SLIOL) and the dorsal intercarpal ligament (DIC) which they termed the Dorsal Capsulo-ligamentous Scapholunate Septum (DCSS). (Figure 1)



Figure 1 The Dorsal Capsulo-ligamentous Scapholunate Septum (DCSS) is thought to be an important stabilizer of the SL joint, which may have therapeutic and prognostic implications.

There are other factors that also determine carpal kinematics. For example ligamentous laxity and carpal morphology affect carpal kinematics.¹² The lunate morphology determines the kinematics of the normal scaphoid, and the abnormal scaphoid.¹³⁻¹⁵ Viegas et al⁴¹ classified lunate morphology as either type 1 or 2, according to the number of facets (one or two, respectively) present on the midcarpal surface of the bone. Lunate type is associated with carpal pathology. Type 1 lunate wrists have a higher incidence of Dorsal Intercalated Segment Instability deformity in the setting of scaphoid non union¹⁴, and type 2 lunate wrists are associated with proximal hamate⁴¹ and scapho-trapezial-trapezoidal joint⁴² degeneration. Lunate morphology is associated with differences in the ligamentous anatomy⁴³ and the kinematics of the carpus, particularly in the central carpal column (radius-lunate-capitate)^{13, 15} (Figure 2)⁴⁷.




Figure 2 Type 1 and type 2 lunate shown in these cadaveric wrists. The type 1 has a single distal facet to articulate with the capitate. The type 2 lunate has 2 distal facets that tend to lock the midcarpal joint. (Used with permission from Fogg)⁴⁷




Figure 3
Scaphoid rotation and flexion

There are anatomic differences in the ligament attachments of the scaphoid. The ligament attachments may predispose to rotation or flexion of the scaphoid. (Figure 3 - used with permission from Quentin Fogg) ⁴⁷

The majority of wrist flexion and extension motion occurs at the three radio-carpal articulations of a type 2 lunate wrist, with the midcarpal articulations being comparatively restricted. In contrast, type 1 lunate wrists have greater motion at the luno-capitate joint and less at the radio-lunate joint during the same wrist movement ¹⁵ (Figure 4).




Figure 4 The wrist type determines the dominant and restricted articulations. In the type 1 wrists, the dominant articulation alternates between the radiocarpal and midcarpal joints. In type 2 wrists, all the midcarpal articulations are restricted and all the radiocarpal articulations are dominant. (In plane motion with wrist flexion (15°) and extension (15°). Dashed line = Dominant articulation ($\geq 50\%$). Solid bar = restricted articulation ($< 4^\circ$). (Copyright G.I. Bain. Used with permission).

Pathoanatomy

The mechanism of injury is usually a fall onto an outstretched hand. With the wrist in extension there is a risk of sustaining an injury to the scapholunate ligament, or alternatively a fracture of the scaphoid or distal radius can occur. The scapholunate ligament injury can be part of a perilunate injury⁵⁸, or part of a carpal dislocation (Figure 5).




Figure 5 Dislocation classification, the spectrum of radiocarpal dislocations that can occur. These include the greater arc (1), lesser arc (2), intra-lunate arc (3) and the inferior arc (4).⁵⁷

History and Physical Examination

The history reported by the patient with scapholunate dissociation usually includes weakness and pain with strenuous activities.⁷ Physical findings usually include swelling in the radial snuffbox or tenderness over the scapholunate interval just distal to Lister's tubercle, pain at the extremes of wrist extension and especially radial deviation, and a positive ballottement test (dorsal volar stress manipulation of the scapholunate interval). Subluxation of the proximal pole of the scaphoid associated with a clunk during dynamic wrist loading (the Watson scaphoid shift test) frequently is present on dynamic testing.¹⁶ The examiner's thumb applies pressure to the scaphoid tubercle as the patient's wrist is brought from a position of ulnar deviation and slight extension to radial deviation and slight flexion. The scaphoid will normally flex and pronate during this manoeuvre, but in scaphoid instability the manoeuvre will be painful, and thumb pressure will force the proximal scaphoid from the scaphoid fossa onto the dorsal articular lip of the radius. Relief of thumb pressure allows the scaphoid proximal pole to spontaneously reduce, often with an audible or palpable "clunk." Patients with an appropriate history and a positive scaphoid shift test should be considered as having a suspected SLIL disruption and should be evaluated further with appropriate imaging or arthroscopy.




Figure 6 Watson's test: The examiner's thumb applies pressure to the scaphoid tubercle as the patient's wrist is brought from a position of ulnar deviation and slight extension to radial deviation and slight flexion. Assess for pain and click or clunk, due to reduction of dorsal scaphoid subluxation.

Imaging

Assessment of the unstable wrist includes plain radiology in all cases, and advanced imaging is often required to determine staging and as part of pre-operative planning.

Plain Radiographs

A complete radiographic assessment with six views of the wrist (postero-anterior, lateral, radial deviation, ulnar deviation, flexion, and extension) is performed. In a patient with scapholunate dissociation, standard PA view (neutral radioulnar deviation) shows an increased scapholunate gap (≥ 3 mm compared with the opposite wrist), and the cortical ring sign of the flexed scaphoid. Lateral radiographs best show scaphoid flexion and lunate extension relative to the radius. SL dissociation should be suspected if the scapholunate angle is greater than the normal 45° to 60° (DISI pattern) (Figure 7). Plain radiographs can be used to identify associated injuries, including the degeneration, which usually begins at the radial styloid, and later can involve the midcarpal joint (Figure 8).




Figure 7 SLA The scapholunate angle: Identify the volar and dorsal distal cusps of the lunate. Draw a line joining these 2 points, which is the alignment of the lunate. The scapholunate angle is the angle between a line drawn perpendicular to the alignment of the lunate and the line along the volar aspect of the scaphoid. Normal is $30^\circ - 60^\circ$. Mean is 47° .



Figure 8 The plain radiographs demonstrate the natural progression of SLAC wrist with degeneration at the radial styloid, then the midcarpal joint.

Cineradiography

Even in static SLD, in which the diagnosis can be made on standard radiographs, obtaining further information using cineradiography is useful. Cineradiography shows not only abnormal movement between the scaphoid and lunate, but also substantial changes in the movement of the midcarpal joint. The hamate-triquetrum relationship normally changes from full engagement in ulnar deviation to complete disengagement in radial deviation; in SLD patients with DISI, this joint remains permanently engaged.

Arthrogram

Contrast is sequentially injected in the midcarpal and radiocarpal joints and scans obtained after each injection. These may be useful in further defining partial tears of the scapholunate ligaments, and in discovering other local problems, such as osteochondral defects or capsular ligament ruptures.¹⁷ When interpreting these scans, care must be taken not to confuse degenerative perforations, or anatomic variants of the scapholunate membrane with true ligament ruptures (Figure 9)^{41, 54}. However, there are several limitations to arthrography, and its use has diminished substantially in favor of arthroscopy.




Figure 9 CT arthrogram of wrist with contrast within the SL interval.

MRI

MRI provides an assessment of the scapholunate ligament integrity, identification of diastasis and chondral changes. The resolution of the scan can make assessment of partial or complete tears unreliable, but improved resolution is certainly much better than previously.^{18, 19} With the MRI the ligament can be assessed, and also the degeneration over the radial styloid (Figure 10).



Figure 10 MRI of the wrist, demonstrating the SL ligament, and adjacent carpus.

Arthroscopy

Wrist arthroscopy is regarded by many authors as the gold standard technique in the diagnosis of intracarpal derangements.¹⁸⁻²³ Three-compartment arthrography will identify perforations of the intercarpal ligaments, but it does not provide accurate localization of the tears or the extent of instability. Arthroscopy has the advantage of direct visualization of the ligaments (Figure 11), and is the most accurate technique for describing the degree of injury of the interosseous ligament, the cartilage, to distinguish fresh from chronic lesions and to analyse concomitant injuries to other structures.²⁴




Figure 11 Normal scapholunate ligament scope (a) Normal scapholunate ligament in the radiocarpal joint, with the ligament of Testut in the background. (b) The probe can be used to palpate the ligament and ensure a concealed tear is not missed.

The state of the ligament, the extent of the ligament injury, and whether it is a repairable ligament stump can be assessed directly. Associated haemorrhage, synovitis, chondral damage, and degenerative changes (e.g. radial styloid degenerative osteoarthritis) can also be visualized.

When infiltration into the midcarpal joint is being performed, a leakage of saline solution through the radiocarpal portals indicates that there must be a tear of the lunotriquetral ligament or scapholunate ligament. This is the same concept as that seen with an arthrogram where the midcarpal joint is injected and a leakage of contrast is seen in the radiocarpal joint on follow-up radiographs.²⁴

From the midcarpal joint, the degree of laxity between the scapholunate interval can be assessed. Geissler et al. described a classification for assessment of scapholunate instability. The functional significance of the ligament injury can be assessed as well – that is, the presence of a tear with or without associated significant instability (as identified in the midcarpal joint). Under the same anaesthetic, a fluoroscopic assessment of the wrist can be performed. If this is performed before draping, then the opposite wrist can be used for comparison. This examination should include placing the wrist in a neutral position, moving to full ulnar deviation, applying an axial load, and also applying traction across the wrist to determine whether there is abnormal distal translation of the scaphoid.²⁴




Figure 12 Scapholunate ligament tear (a) Tear of the ligament seen from the radiocarpal joint. (b) Diastasis of the scapholunate interval interval, when the Watson’s test is performed under arthroscopic vision²⁴.

The scapholunate instability test of Watson et al. can be performed under fluoroscopy or arthroscopic vision (or both) (Figure 12) ²⁴. Abnormal widening of the scapholunate interval and subluxation or dislocation of the scaphoid over the dorsal rim of the distal radius can be identified.

Lunotriquetral instability can also be assessed by use of the same arthroscopic assessment techniques and specific provocation tests. Pressure is placed directly onto the pisiform and on the dorsal aspect of the lunate. By squeezing the lunate volarly and

the triquetrum dorsally, lunotriquetral instability is identified.²⁴ The wrist is taken through radial and ulnar deviation with direct visualization of the lunotriquetral articulation.

Arthroscopic Classification

Because arthroscopy of the wrist is one of the most accurate means of evaluation of scapholunate instability, the classifications deduced from these findings are widely used. Geissler has proposed a method of quantifying the degree of interosseous ligament injury (Table 2), by probe placement into the scapholunate interval from the radiocarpal and midcarpal joint on wrist arthroscopy. (Figure 13). If the scope can be advanced from the radiocarpal to the midcarpal joint, it is classified as a grade 4 injury (Figure 14).



Figure 13 Probe in the scapholunate interval in the midcarpal joint. It is normal for the probe to not be able to be admitted into the interval.



Figure 14 Grade 4 tear, with the capitate seen in the interval between the scaphoid (left) and lunate (right).

Table 2: Geissler Arthroscopic Grading System ²⁵

| Grade | Description |
|--------------|---|
| I | Attenuation/haemorrhage of SLIL (viewed from radiocarpal space). No midcarpal malalignment |
| II | Attenuation/haemorrhage of SLIL (viewed from radiocarpal space) AND step off / incongruency of carpal alignment. Slight gap between carpals (less than width of probe) |
| III | Step off / incongruency of carpal alignment (viewed from both radiocarpal and midcarpal space) AND SL gap large enough to pass probe between carpals |
| IV | Step off / incongruency of carpal alignment (viewed from both radiocarpal and midcarpal space), gross instability, AND 2.7- mm arthroscope can pass through the gap between the scaphoid and lunate (positive “drive-through sign”) |

Recently, the European Wrist Arthroscopy Society (EWAS) classification for SL dissociation was introduced. This is a more comprehensive classification that includes the site of the scapholunate ligament attenuation or tear.²⁶

References

1. Linscheid RL, Dobyns JH, Beabout JW, Bryan RS. Traumatic instability of the wrist. Diagnosis, classification, and pathomechanics. *The Journal of bone and joint surgery American volume*. 1972 Dec;54(8):1612-32.
2. Taleisnik J. Classification of carpal instability. *Bulletin of the Hospital for Joint Diseases Orthopaedic Institute*. 1984 Fall;44(2):511-31.
3. Cooney WP, Dobyns JH, Linscheid RL. Arthroscopy of the wrist: anatomy and classification of carpal instability. *Arthroscopy : the journal of arthroscopic & related surgery : official publication of the Arthroscopy Association of North America and the International Arthroscopy Association*. 1990;6(2):133-40.
4. Hodge JC, Gilula LA, Larsen CF, Amadio PC. Analysis of carpal instability: II. Clinical applications. *The Journal of hand surgery*. 1995 Sep;20(5):765-76; discussion 77.
5. Larsen CF, Amadio PC, Gilula LA, Hodge JC. Analysis of carpal instability: I. Description of the scheme. *The Journal of hand surgery*. 1995 Sep;20(5):757-64.
6. Cassidy C, Ruby LK. Carpal instability. *Instructional course lectures*. 2003;52:209-20.
7. Watson HK, Weinzweig J, Zeppieri J. The natural progression of scaphoid instability. *Hand clinics*. 1997 Feb;13(1):39-49.
8. Berger RA. The gross and histologic anatomy of the scapholunate interosseous ligament. *The Journal of hand surgery*. 1996 Mar;21(2):170-8.
9. Kitay A, Wolfe SW. Scapholunate instability: current concepts in diagnosis and management. *The Journal of hand surgery*. 2012 Oct;37(10):2175-96.
10. Elsaidi GA, Ruch DS, Kuzma GR, Smith BP. Dorsal wrist ligament insertions stabilize the scapholunate interval: cadaver study. *Clinical orthopaedics and related research*. 2004 Aug(425):152-7.
11. Overstraeten LV, Camus EJ, Wahegaonkar A, Messina J, Tandara AA, Binder AC, et al. Anatomical Description of the Dorsal Capsulo-Scapholunate Septum (DCSS)- Arthroscopic Staging of Scapholunate Instability after DCSS Sectioning. *Journal of wrist surgery*. 2013 May;2(2):149-54.
12. Garcia-Elias M, Ribe M, Rodriguez J, Cots M, Casas J. Influence of joint laxity on scaphoid kinematics. *Journal of hand surgery*. 1995 Jun;20(3):379-82.
13. Galley I, Bain GI, McLean JM. Influence of lunate type on scaphoid kinematics. *The Journal of hand surgery*. 2007 Jul-Aug;32(6):842-7.
14. Haase SC, Berger RA, Shin AY. Association between lunate morphology and carpal collapse patterns in scaphoid nonunions. *The Journal of hand surgery*. 2007 Sep;32(7):1009-12.
15. Bain G, et al. The Effect Of Lunate Morphology on Three-Dimensional Carpal Kinematics *J Hand Surg Am* 2014

16. Watson HK, Ashmead Dt, Makhlof MV. Examination of the scaphoid. *The Journal of hand surgery*. 1988 Sep;13(5):657-60.
17. Theumann N, Favarger N, Schnyder P, Meuli R. Wrist ligament injuries: value of post-arthrography computed tomography. *Skeletal radiology*. 2001 Feb;30(2):88-93.
18. Mahmood A, Fountain J, Vasireddy N, Waseem M. Wrist MRI Arthrogram v Wrist Arthroscopy: What are we Finding? *The open orthopaedics journal*. 2012;6:194-8.
19. Magee T. Comparison of 3-T MRI and arthroscopy of intrinsic wrist ligament and TFCC tears. *AJR American journal of roentgenology*. 2009 Jan;192(1):80-5.
20. Binder AC, Kerfant N, Wahegaonkar AL, Tandara AA, Mathoulin CL. Dorsal wrist capsular tears in association with scapholunate instability: results of an arthroscopic dorsal capsuloplasty. *Journal of wrist surgery*. 2013 May;2(2):160-7.
21. Katschnig I, Prosquill E. [Arthroscopy of the wrist: compared results of MRT and arthrography and outcome in the arthroscopy -- an examination 1998 to 2003]. *Handchirurgie, Mikrochirurgie, plastische Chirurgie : Organ der Deutschsprachigen Arbeitsgemeinschaft fur Handchirurgie : Organ der Deutschsprachigen Arbeitsgemeinschaft fur Mikrochirurgie der Peripheren Nerven und Gefasse* 2006 Apr;38(2):104-8.
22. Kozin SH. The role of arthroscopy in scapholunate instability. *Hand clinics*. 1999 Aug;15(3):435-44, viii.
23. Whipple TL. The role of arthroscopy in the treatment of scapholunate instability. *Hand clinics*. 1995 Feb;11(1):37-40.
24. Bain GI, Munt J, Turner PC. New advances in wrist arthroscopy. *Arthroscopy : the journal of arthroscopic & related surgery : official publication of the Arthroscopy Association of North America and the International Arthroscopy Association*. 2008 Mar;24(3):355-67.
25. Geissler WB, Freeland AE, Savoie FH, McIntyre LW, Whipple TL. Intracarpal soft-tissue lesions associated with an intra-articular fracture of the distal end of the radius. *The Journal of bone and joint surgery American volume*. 1996 Mar;78(3):357-65.
26. Messina JC, Van Overstraeten L, Luchetti R, Fairplay T, Mathoulin CL. The EWAS Classification of Scapholunate Tears: An Anatomical Arthroscopic Study. *Journal of wrist surgery*. 2013 May;2(2):105-9.
27. Garcia-Elias M, Lluch AL, Stanley JK. Three-ligament tenodesis for the treatment of scapholunate dissociation: indications and surgical technique. *The Journal of hand surgery*. 2006 Jan;31(1):125-34.
28. Hagert E, Garcia-Elias M, Forsgren S, Ljung BO. Immunohistochemical analysis of wrist ligament innervation in relation to their structural composition. *The Journal of hand surgery*. 2007 Jan;32(1):30-6.
29. Salva-Coll G, Garcia-Elias M, Hagert E. Scapholunate instability: proprioception and neuromuscular control. *Journal of wrist surgery*. 2013 May;2(2):136-40.

30. Salva-Coll G, Garcia-Elias M, Llusa-Perez M, Rodriguez-Baeza A. The role of the flexor carpi radialis muscle in scapholunate instability. *The Journal of hand surgery*. 2011 Jan;36(1):31-6.
31. Melone CP, Jr., Polatsch DB, Flink G, Horak B, Beldner S. Scapholunate interosseous ligament disruption in professional basketball players: treatment by direct repair and dorsal ligamentoplasty. *Hand clinics*. 2012 Aug;28(3):253-60, vii.
32. Stuffmann ES, McAdams TR, Shah RP, Yao J. Arthroscopic repair of the scapholunate interosseous ligament. *Techniques in hand & upper extremity surgery*. 2010 Dec;14(4):204-8.
33. Short WH, Werner FW, Sutton LG. Dynamic biomechanical evaluation of the dorsal intercarpal ligament repair for scapholunate instability. *The Journal of hand surgery*. 2009 Apr;34(4):652-9.
34. Szabo RM. Scapholunate ligament repair with capsulodesis reinforcement. *The Journal of hand surgery*. 2008 Nov;33(9):1645-54.
35. Bleuler P, Shafiqhi M, Donati OF, Gurunluoglu R, Constantinescu MA. Dynamic repair of scapholunate dissociation with dorsal extensor carpi radialis longus tenodesis. *The Journal of hand surgery*. 2008 Feb;33(2):281-4.
36. Wahegaonkar AL, Mathoulin CL. Arthroscopic dorsal capsulo-ligamentous repair in the treatment of chronic scapho-lunate ligament tears. *Journal of wrist surgery*. 2013 May;2(2):141-8.
37. Mathoulin CL, Dauphin N, Wahegaonkar AL. Arthroscopic dorsal capsuloligamentous repair in chronic scapholunate ligament tears. *Hand clinics*. 2011 Nov;27(4):563-72, xi.
38. Brunelli F, Spalvieri C, Bremner-Smith A, Papalia I, Pivato G. [Dynamic correction of static scapholunate instability using an active tendon transfer of extensor brevis carpi radialis: preliminary report]. *Chirurgia de la main*. 2004 Oct;23(5):249-53.
39. Surdziel P, Romanowski L, Czarnecki P, Kaczmarek L, Splawski R. Dynamic correction of scapholunate instability using ECRB tendon transfer. *Ortopedia, traumatologia, rehabilitacja*. 2006 Apr 28;8(2):134-8.
40. Rosenwasser MP, Miyasajsa KC, Strauch RJ. The RASL procedure: reduction and association of the scaphoid and lunate using the Herbert screw. *Techniques in hand & upper extremity surgery*. 1997 Dec;1(4):263-72.
41. Viegas SF, Wagner K, Patterson R, Peterson P. Medial (hamate) facet of the lunate. *J Hand Surg Am*. Elsevier; 1990;15(4):564-71.
42. McLean J, Turner, P.C., Bain GI, Rezaian N, Field J, Fogg Q. An association between lunate morphology and scaphoid-trapezium-trapezoid arthritis. *J Hand Surg Eur Vol*. 2009 Dec;34(6):778-82.
43. Nakamura K, Patterson RM, Moritomo H, Viegas SF. Type I versus type II lunates: Ligament anatomy and presence of arthrosis. *J Hand Surg Am*. 2001 May;26(3):428-36.

44. Bain GI, Watt AC. The outcome of scaphoid excision and four corner arthrodesis for advanced carpal collapse at a minimum of ten years. *Journal of Hand Surgery (American)*, 2010, 35A:719-725.
45. Van Riet RP, Bain GI. Three corner wrist fusion using memory staples. *Techniques in Hand and Upper Extremity Surgery*. 10(4):259-262, 2006.
46. Singh HP, Dias JJ, Phadnis J, Bain GI. Comparison of the Clinical and Functional Outcomes Following 3- and 4- Corner Fusions. *JHS (Am)* (40), 6, 1117-1123. 2015.
47. Fogg Q. Scaphoid Variation and an Anatomical Basis for Variable Carpal Mechanics. PhD Thesis. Department of Anatomy, University of Adelaide. 2000-2003.
48. Watts AC, McLean JM, Fogg Q, Bain GI. Scaphoid Anatomy In: The Scaphoid. Editor Slutsky DJ and Slade JF III. New York: Thieme, 2011:3-10.
49. Bain GI, Sood A, Yeo CJ. RSL Fusion with Excision of Distal Scaphoid and Triquetrum: A Cadaveric Study. *JWS* 2014; 03(01): 037-041, doi: 10.1055/s-0033-1364095.
50. Vance MC, Hernandez JD, Didonna ML, Stern PJ. Complications and outcome of four-corner arthrodesis: Circular plate fixation versus traditional techniques. *J Hand Surg AM* 2005;30:1122-7.
51. Trail IA, Murali R, Stanley JK, Hayton MJ, Talwalkar S, Sreekumar R, Birch A. The long-term outcome of four-corner fusion. *Journal of Wrist Surgery*. 2015;4(2):128-133.
52. Wilhelm A. Partial joint denervation: wrist, shoulder and elbow. *Plast Reconstr Surg*. 2010; Jul; 126(1):345-7.
53. Chim H, Moran S. Long-Term Outcomes of Proximal Row Carpectomy: A Systematic Review of the Literature. *J Wrist Surg*. 2012 Nov; 1(2):141-148.
54. Boyer JS, Adams BD. Distal radius hemiarthroplasty combined with proximal row carpectomy: case report. *Iowa Orthop J*. 2010, 30:168-173.
55. Hastings H, Weiss AP, Quenzer D, Wiedeman GP, Hanington KR, Strickland JW. Arthrodesis of the wrist for post-traumatic disorders. *J Bone J Surg AM*. 1996; 78:897-902.
56. Ross M, Loveridge J, Cutbush K, Couzens G. Scapholunate Ligament Reconstruction. *J Wrist Surg*. 2013 May; 2(2):110-115.
57. Bain GI, McGuire DT. Decision Making for Partial Carpal Fusions. *Journal of Wrist Surgery*. 2012;01(02):103-114
58. Herzberg G, Comtet JJ, Linscheid RL, Amadio PC, Conney WP, Stadler J. Perilunate dislocations and fracture-dislocations: a multicenter study. *J Hand Surg AM* 1993;18(5):768-779.
59. Brunelli GA, Brunelli GR. A new surgical technique for carpal instability with scapholunate dissociation. *Surg Technol Int*. 1996;5:370-374.
60. Bain GI, Watts AC, McLean J, Lee YC, Eng K. Cable-augmented, Quad Liagment

Tenodesis Scapholunate Reconstruction. J Wrist Surgery. 2015:

61. Chee KG, Chin AYH, Chew EM, Garcia-Elias M. Antipronation spiral tenodesis - a surgical technique for the treatment of perilunate instability. *The Journal of Hand Surgery*. 2012;Volume 37, p 2611-18.
62. Lee SK, Zlotolow DA, Sapienza A, Karia R, Yeo J. Biomechanical comparison of 3 methods of scapholunate ligament reconstruction. *The Journal of Hand*, 2014;(39) p643-650.
63. Cha SM, Shin HD, Kim KC. Clinical and radiological outcomes of scaphoidectomy and 4-corner fusion in scapholunate advanced collapse at 5 and 10 years. *Ann Plast Surg*. 2013 Aug;71(2):166-9.
64. Krukhaug Y, Lie SA, Havelin LI, Furness O, Hove LM. Results of 189 wrist replacement. A report from the Norwegian Arthroplasty Register. *Acta Orthop*. 2011; Aug;82(4):405-9. doi:10.3109/17453674.2011.588858. Epub.2011.